

The Relation Between Body Mass Index (BMI) and Bronchial Asthma Control in Children in Tabuk City - 2017

Meshari Faisal Alenezi¹, Saleh Hammad Alhawiti^{1*}, Mohammed Ahmed Alhejaili¹, Adulleelah Mohammed Alqarni¹, Khalid Awd Albalawi¹, Tariq H Merhani²

¹Sixth Year Medical Student, Faculty of Medicine, University of Tabuk, Tabuk, KSA.

²Associate Professor, Faculty of Medicine, University of Tabuk, Tabuk, KSA.

ABSTRACT

Introduction: Most Saudi asthmatic children/adolescents visiting the ED had poor control of symptoms; indeed, none achieved complete control, which is related to deficient medication compliance and improper medication inhaler device use; deficient knowledge about asthma was also another factor hindering control. The objective from this study To analyze the relationship between obesity (measured by body mass index) and asthma control in the children known to have bronchial asthma.

Method: A cross-sectional study was carried out in ten primary and intermediate schools in Tabuk city, Saudi Arabia. A self-administered questionnaire for parents with asthmatic children was sent with each school pupil, to be filled by one of his parents at home and then to be returned on the next day. The questionnaire responses were analyzed using the Statistical Package for the Social Science (SPSS Inc. Chicago, IL, USA) version 20.

Result: 68% of the parents education level was intermediate and secondary school. Obese children with uncontrolled asthma were higher than obese children with controlled asthma. (33.3%).(P<.05). 23.1% of children with uneducated parents have uncontrolled bronchial asthma, compared to other children with primary, intermediate, secondary, and graduated parents. (P<0.05). Dust triggers bronchial asthma in 80% of the patients either alone or coexisted with other triggering factors. Only (11%) of asthmatic children use peak

flow meter while the majority of them, which is about (61%), did not use it. Furthermore, about (28%) of parents with asthmatic children did not know this meter.

Conclusion: Abnormal BMI associated with poor control of bronchial asthma among children, the higher the BMI, the poorer control of bronchial asthma, at the same time, underweight BMI children will suffer of poor control of bronchial asthma. Parental education proportionally associated with bronchial asthma control. We need more efforts among pediatricians and physicians to increase awareness of importance and use of Peak Flow Meter.

Key words: Bronchial Asthma, Control, Body Mass Index, Childhood.


*Correspondence to:

Saleh Hammad Alhawiti,
Sixth Year Medical Student,
Faculty of Medicine, University of Tabuk, Tabuk, KSA.

Article History:

Received: 27-08-2017, Revised: 28-09-2017, Accepted: 24-10-2017

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2017.3.6.022	

INTRODUCTION

Bronchial asthma is a heterogeneous disorder of the conducting airways involving chronic airway inflammation, declining airway function and tissue remodeling.¹ Asthma is a condition whereby one's airways can narrow and produce extra mucus causing symptoms of coughing, wheezing and shortness of breath.² The prevalence of asthma has rapidly increased over the last few decades to epidemic proportions and there are now an estimated 300 million individuals of all ages, ethnic groups and countries suffering from bronchial asthma, a total that is expected to rise dramatically over the next 15–20 year.¹ It is one of the most common chronic diseases in Saudi Arabia, affecting more than 2 million Saudis.³⁻⁵ Its impact is manifested in patients, their families, and the community as a whole in terms of lost work and school

days, poor quality of life, frequent emergency department (ED) visits, hospitalizations, and deaths. Asthma is associated with enormous healthcare expenditures and, despite the advances in effective therapy, the consequential economic burden associated with disease control and morbidity continues to escalate.⁶ It is estimated that around 250,000 people die prematurely each year as a result of asthma. Concepts of asthma severity and control are important in evaluating patients and their response to treatment.¹ There is increasing and convincing epidemiological evidence of an association between obesity and asthma. Cross-sectional studies almost uniformly show an increased prevalence of asthma in obese subjects as compared with their leaner counterparts, both in adults and in children.⁷ An even more important question is why

asthma symptoms in obese patients are often so difficult to control.⁸ Obese patients with asthma are indeed less responsive to standard asthma therapy and remain uncontrolled even when prescribed high doses of inhaled corticosteroids.⁹ Weight gain and obesity have been associated with many chronic conditions like diabetes, hypertension and sleep apnea. Asthma can now be added to this list of chronic conditions affected by obesity. An increase in asthma has been seen in both children and adults with obesity, particularly among women. Obesity itself is now listed as a risk factor for the development of asthma. A person who is affected by obesity has a higher chance of developing asthma that responds less well to medications. A clear-cut relationship has also been described whereby the higher one's weight, the higher one's chances of developing asthma. Obesity has specifically been associated with an increase in daily asthma symptoms, missed workdays, an increased use of bronchodilator medications and an increased risk of hospitalization.² There is a relationship between the trend towards obesity and asthma. Moreover, it has been reported that asthma symptoms are associated with obesity.¹⁰ Different hypothesis has been proposed regarding this relationship between obesity and asthma, one hypothesis relies on the facts that obesity leads to a decrease in lung volumes including the tidal volume which may increase airway obstruction, adipose tissue derived hormones and cytokines such as leptin, adiponectin, tumor necrosis factor and interleukins that play an inflammatory role.¹¹ Adipose tissue produces a number of mediators, termed adipokines, which have significant metabolic effects. One of these adipokines, adiponectin, is actually decreased in the obese. primary metabolic effects of adiponectin are on glucose regulation and fatty acid metabolism, adiponectin is also anti-inflammatory.^{12,13}

The normocaloric dietary intervention was associated with improved Asthma Related Quality of Life and some asthma control. Dietary programs may serve as a complementary non-pharmacologic therapeutic strategy in obese adolescents who have asthma.¹⁴ Dietary intervention can induce acute weight loss in obese asthmatic children, with improvements in static lung function, asthma control, and self-reported quality of life. Systemic and airway inflammation did not change after weight loss.¹⁴ Obese children were more symptomatic than non-obese children were. Obese children reported more cough and wheezing than non-obese children. Asthma, as diagnosed by physician and inhaler use, was more common among obese children than non-obese children.¹⁵ On the other hand, childhood obesity is one of the most

serious public health challenges of the 21st century. The problem is global and is steadily affecting many low- and middle-income countries, particularly in urban settings. The prevalence has increased at an alarming rate.¹⁶ Globally, in 2015 the number of overweight children under the age of five, is estimated to be over 42 million, at the same time, overweight and obese children are likely to stay obese into adulthood. Overweight and obesity, as well as their related diseases, are largely preventable. Prevention of childhood obesity therefore needs high priority.¹⁷ Saudi Arabia the National Growth Study showed that the overall prevalence of obesity in children and adolescents from 5 to 18 years was 11.3%.^{18,19}

METHODS

We have conducted a descriptive cross-sectional study in twelve primary and intermediate schools, in Tabuk city, Saudi Arabia. The study was conducted during the period from January to May 2017. The schools were selected by random sampling. Sampling was stratified for the different geographical areas of the city. Our study included children with bronchial asthma aged 5-16 years old.

We recruited 400 child with bronchial asthma, and a self-administered questionnaire, previously validated,²⁰ require information about: (symptom of bronchial asthma regarding severity and control, and most common causes of exacerbations) was sent with each pupil to his parents, to be filled by one of them and then to be returned back on the next day. A letter that explains the objectives of the study and asks for parental consent was sent with the questionnaire. Each pupil's weight and height measured by medical professional and well trained students, then Body Mass Index (BMI) was calculated as $\text{Weight} \backslash \text{Height}^2$. BMI classified to Underweight, Normal, Overweight and Obese according to Saudi growth charts of BMI.

The questionnaire responses were analyzed using the Statistical Package for the Social Science (SPSS Inc. Chicago, IL, USA) version 20. Categorical variables were described by frequencies and percentages. Descriptive analysis involving Chi-square test was used to test significance of association between categorical variables. The level of significance was set at $P < 0.05$. The research was approved by the local Research Committee of the Faculty of Medicine, University of Tabuk. Official letters were sent to the selected primary and intermediate schools. Parents of the pupils were asked to give their written consents before participation in the study.

Table-1: General characteristics of the participants (n= 400)

Character		
Age	Mean (SD) (y)	9.70 (2.81)
	Range (y)	5 – 14
Gender	Male (n (%))	274 (68.5%)
	Female (n (%))	126 (31.5%)
Education	Not educated (n (%))	052 (13%)
	Primary/Intermediate/ Secondary (n (%))	292 (73%)
	Graduate (n (%))	056 (14%)
Income	Poor (n (%))	068 (17%)
	Average (n (%))	308 (77%)
	High (n (%))	024 (6%)

Table-2: Bronchial asthma control in relation to BMI

BMI	Control n(%)	Partial control n(%)	Uncontrolled n(%)	Total n(%)
Underweight	64(43.25%)	64(43.25%)	20(13.5%)	148(100%)
Normal	76(51.4%)	64(43.2%)	8(5.4%)	148(100%)
Overweight	32(57.1%)	12(21.4%)	12(21.4%)	56(100%)
Obese	24(50.0%)	8(16.7%)	16(33.3%)	48(100%)
Total N(%)	196(100%)	148(100%)	56(100%)	400(100%)

Table-3: Bronchial asthma control in relation to education of parents

Education	Controlled n(%)	Partially controlled n(%)	Uncontrolled n(%)	Total n(%)
Uneducated	24(46.1%)	16(30.8%)	12(23.1%)	52(100%)
Primary\ Intermediate\ Secondary School	152(52%)	104(35.6%)	36(12.4%)	292(100%)
Graduated	20(35.7%)	28(50%)	8(14.3%)	56(100%)
Total n(%)	196(100%)	148(100%)	56(100%)	400(100%)

Table-4: Prevalence of triggering factors in the study population (n=400)

Triggering Factors	(n)	%
Dust	320	80%
Common Cold	196	49%
Cigarette Smoking	184	46%
Cold Weather	172	43%
Physical Activity	88	22%
Pits	64	16%
Pollens	28	7%

Table-5: Knowledge and use of Peak Flow Meter

Is the asthmatic Px using Peak Flow Meter	n=400	%
Yes	44	11
No	244	61
I do not know it	112	28
Total	400	100

RESULTS

Table-1 shows general characteristics of the participants and their parents. Age of participants ranged from 5 to 14 year old, with a mean (SD) of 9.70 (2.81), the majority were male (68.5%), and (68%) of the parents education level was intermediate and secondary school.

Table-2 shows significant relation between bronchial asthma control and BMI ($P < 0.05$). Of total 400 participants, (37%) were under weight, (37%) were normal, (14%) were overweight, (12%) were obese. within underweight participants, (43.2%) of them were controlled, (43.2%) were partially controlled, (13%) were uncontrolled. For normal BMI children, (51.4%) of them were good control, (43.2%) were partially controlled, (5.4%) were uncontrolled. For overweight children, (57.1%) of them were good control, (21.45%) were partially controlled, (21.45%) were uncontrolled. For obese children (50%) of them were control, (16.7%) were partially controlled, (33.3%) were uncontrolled.

Obese children with uncontrolled bronchial asthma were higher than obese children with controlled bronchial asthma. While normal BMI children with controlled bronchial asthma higher than normal BMI children with uncontrolled bronchial asthma. overweight children with uncontrolled asthma higher than overweight children with controlled asthma. Under weight children with controlled and uncontrolled bronchial asthma were nearly equal. Table-3 shows significant relation between bronchial asthma control and level of parents education ($p < .05$). (23.1%) of children with uneducated parents have uncontrolled bronchial asthma, compared to other children with primary, intermediate, secondary, and graduated parents (00.0%), (10%), (15.8%), and (14.3) respectively. No significant relation between bronchial asthma control in child and gender.

Table-4 shows the prevalence of triggering factors in the study population. Dust triggers bronchial asthma in 80% of the patients either alone or coexisted with other triggering factors. As same as for common cold (49%) cigarette smoking (46%) and cold weather (43%). Table-5 shows the use and knowledge of Peak Flow Meter. Asthmatic children who use peak flow meter were (11%), children who do not use it were (61%), and children who did not know about it were (28%).

DISCUSSION

Asthma in children is a leading cause of emergency department visits, hospitalizations and missed school days. Unfortunately, childhood asthma can't be cured, and symptoms may continue into adulthood. But with the right treatment, you and your child can keep symptoms under control and prevent damage to growing lungs.²¹ In this study, uncontrolled bronchial asthma measured by GINA guide lines of bronchial asthma severity and control. Based on our finding, Obese children have the highest rate of uncontrolled bronchial asthma (33.3%) compared to overweight, and underweight children who have uncontrolled bronchial asthma were (21.4%), and (13.5%) respectively. Children with normal BMI with uncontrolled bronchial asthma were (5.4%). Similar finding were reported in other countries suggest that childhood obesity is associated with increased risk of worse bronchial asthma control and exacerbation.²² Other similar finding that Results identify

higher BMI and obesity as potential behavioral factors related to worse asthma control and quality of life, but not asthma severity, and suggest important avenues for asthma management and control initiatives.²³ This study showed that, obese, overweight, and underweight children associated with more uncontrolled bronchial asthma, in both genders, but more in female than in male. Another study showed that higher BMI was associated with more symptom days and exacerbations among females only.²⁴ Other study suggest that there is no statistically significant association between asthma control and BMI.²⁵ In this study, the majority of parents education level was intermediate and secondary school(68%), graduated parents and parents with basic education have less children with uncontrolled bronchial asthma compared to uneducated parents who have more children with uncontrolled bronchial asthma. A study done among children diagnosed with bronchial asthma, demonstrated that children from families with lower education used fewer controller medications than those whose parents were college graduates.²⁶ The prevalence of triggering factors also explored, the most common triggering factors were dust, common cold. A study done in Tabuk city showed that the most common triggering factors of bronchial asthma exacerbation were URTI, and dust.²⁷ It is important to be aware of the common triggering factors in our area to help to control asthma symptoms and decrease the hospitalization by advising patients about the needed techniques for reducing any allergen exposure. Given the increasing morbidity and mortality of asthma, it is clear that therapeutic efforts should be focused on identifying relevant allergens and advising patients about techniques for reducing exposure.²⁸ Measuring peak flow rate in asthma episodes helps to determine the severity of exacerbations and assists in guiding therapeutic decisions in the home, school, practitioner's office, and emergency department.²⁹ Based on the findings in this study, only (11%) of asthmatic children use peak flow meter while the majority of them, which is about (61%), did not use it, and furthermore, about (28%) of parents with asthmatic children did not know this meter.

In conclusion, the majority of children with uncontrolled bronchial asthma were obese children. The most common triggering factors of bronchial asthma in Tabuk city were Dust followed by common cold. The majority of children whom parents were uneducated have uncontrolled bronchial asthma. As the majority of parents for children with bronchial asthma do not use PFM or do not know about it, we need more efforts to increase the awareness about PFM among parent for asthmatic children.

REFERENCES

- World Health Organization. Global surveillance, prevention and control of chronic respiratory diseases: a comprehensive approach, 2007. Accessed at 2017 May 13. Available at <http://www.who.int/gard/publications/GARD%20Book%202007.pdf>
- <http://www.obesityaction.org/educational-resources/resource-articles-2/obesity-related-diseases/obesity-and-asthma>. Accessed at May 14 2017.
- Stewart WF, Ricci JA, Chee E, Morganstein D. Lost productive work time costs from health conditions in the United States: Results from the American Productivity Audit. *J Occup Environ Med* 2003;45:1234-46.
- Al-Mobeireek A. Prescribing for asthmatic children in primary care. Are we following the guidelines? *Saudi Med J* 2003;24:1274.
- Dashash NA, Mukhtar SH. Prescribing for asthmatic children in primary care. Are we following guidelines?*SaudiMedJ* 2003;24:507-11.
- Murdoch JR, Lloyd CM. Chronic inflammation and asthma. *Mutation Research*. 2010 Aug 7; 690(1-2): 24–39.
- Ford ES. The epidemiology of obesity and asthma. *J Allergy Clin Immunol* 2005;115:897–909.
- Quinto KB, Zuraw BL, Poon KY, Chen W, Schatz M, Christiansen SC. The association of obesity and asthma severity and control in children. *J Allergy Clin Immunol* 2011;128:964–969.
- Sutherland ER, Goleva E, Strand M, Beuther DA, Leung DY. Body mass and glucocorticoid response in asthma. *Am J Respir Crit Care Med* 2008;178:682–687.
- Tai A, Volkmer R, Burton A. Association between asthma symptoms and obesity in preschool (4-5 year old) children. *J Asthma* 2009; 46: 362-5.
- Shore SA. Obesity and asthma: lessons from animal models. *J Appl Physiol* 2007; 102: 516-28.
- Dixon AE. Adipokines and asthma. *Chest* 2009; 135:255–256.
- Shore SA, Terry RD, Flynt L, Xu A, Hug Ch. Adiponectin attenuates allergen induced airway inflammation and hyper responsiveness in mice. *J Allergy Clin Immunol* 2006;118:389–95.
- Jensen ME, Gibson PG, Collins CE, Hilton JM, Wood LG. Diet-induced weight loss in obese children with asthma: a randomized controlled trial. *Clin Exp Allergy*.2013Jul;43(7):775-84.
- Bibi H, Shoseyov D et al. The relationship between asthma and obesity in children: is it real or a case of over diagnosis?.*J Asthma*. 2004 Jun;41(4):403-10.
- Obesity: Preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser*; 2000. p. 100-42. Accessed at 2017 May 14. Available on file:///C:/Users/HP/Downloads/WHO_TRS_894%20(4).pdf.
- WHO. Global Strategy on Diet, Physical Activity and Health. Childhood overweight and obesity. Accessed on 2017 May 14. Available at: <http://www.who.int/dietphysicalactivity/childhood/en/>
- Musaiger AO. Overweight and obesity in Eastern mediterranean region: Prevalence and possible causes. *J Obes* 2011;2011:2-17.
- El-Mouzan MI, Foster PJ et al. Prevalence of overweight and obesity in Saudi children and adolescents. *Ann Saudi Med* 2010;30:203-8.
- Pocket guide for asthma management and prevention(for adults and children alder than five years). 2017.Gina Initiative For Asthma.
- Huffaker MF, et al. Pediatric asthma: Guidelines-based care, omalizumab, and other potential biologic agents. *Immunology Allergy Clinics of North America*. 2015;35:129.
- Kenneth B, Bruce L et al. The association of obesity and asthma severity and control in children. *Journal of Allergy and Clinical Immunology*; 128(5), 2011, 964-969.
- Kim L, Simon L et al. Higher BMI is associated with worse asthma control and quality of life but not asthma severity. *Respiratory Medicine*;100(4), April 2006, 648-657.
- Meyer Kattan, M.D., C.M., Rajesh Kumar, M.D. et al. Asthma control, adiposity, and adipokines among inner-city adolescents *J Allergy Clin Immunol.* 2010 March; 125(3): 584–592
- Fariba A, Susanne V et al. Childhood obesity in relation to poor asthma control and exacerbations. *European Respiratory Journal* 2016, 48 (4): 1063-1073.

26. Gong T, Lundholm C et al. Parental Socioeconomic Status, Childhood Asthma and Medication Use – A Population-Based Study. PLoS One. 2014 Sep 4;9(9)
27. Abdalla A, Osama S et al. Basic Research Journal of Medicine and Clinical Sciences. Association of triggering factors with asthma exacerbations among the pediatric population in Tabuk, Kingdom of Saudi Arabia. 2016 September; 5(7):114-118.
28. Platts-M, Ward J et al. Epidemiology of the Relationship between Exposure to Indoor Allergens and Asthma. Int Arch Allergy Immunol 1991;94:339–345.
29. PFM10-National Asthma Education and Prevention Program. Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma; Full Report 2007. Department of Health and Human Services, National Institutes of Health, National Heart, Lung and Blood Institute. 2007. Available on <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.htm>

Source of Support: Nil. **Conflict of Interest:** None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Meshari Faisal Alenezi, Saleh Hammad Alhawiti, Mohammed Ahmed Alhejaili, Adulleelah Mohammed Alqarni, Khalid Awd Albalawi, Tariq H Merhani. The Relation Between Body Mass Index (BMI) and Bronchial Asthma Control in Children in Tabuk City - 2017. Int J Med Res Prof. 2017 Nov; 3(6):108-12. DOI:10.21276/ijmrp.2017.3.6.022