

Knowledge and Attitude Regarding First Aid Management of Epistaxis Among Students of Medical Colleges at King Khalid University, Abha City, Saudi Arabia

Muhannad A. Asiri^{1*}, Sarah M. Mujarri², Shahad S. Alkhashrami², Amal A. Alghamdi², Khalid A. Almalwi², Saeed M. Asiri²

¹Intern, ²Medical Student, College of Medicine, King Khalid University, Abha, Saudi Arabia.

ABSTRACT

Background: Epistaxis is the most common ear, nose and throat complaint presenting to an emergency departments worldwide which is defined as a bleeding from inside the nose or nasal cavity. A vast majority of these patients settle with simple standard first aid measures.

Objective: The aim of this study was to evaluate knowledge and attitude regarding first aid management of epistaxis among health related specialties students at King Khalid University, Abha city, Saudi Arabia.

Methods: A descriptive cross sectional study was undertaken among medical specialties students at King Khalid University, Abha city, Saudi Arabia. Data were collected using an online questionnaire tool, with sum of 24 questions.

Results: Data was collected from 314 medical specialties students between January and March 2017 by using questionnaires which were filled electronically. 205 (65.3%) were male and 109 (34.7%) were female. Their age ranged between 18 and 26 years with a mean of 21.71 ± 1.56 years. Only 131 (41.7%) had received a first aid provider certificate. The medical students were the most respondents (52.2%) while (47.8%) were other medical specialties. Majority of the respondents were fourth year students. 167(83.1%) respondents said that a nosebleed cannot be stopped after 10 to 20 minutes of direct pressure is one of the commonest cause to seek for emergency care. The commonest first aid measures reported to be known by respondents when the patients get shocked, were Pinching the nose (86.3%), Nasal packing (78.9%), Putting the patient in supine position with the head lowered (59.8%), giving Anti shock treatment (56.6%) and Putting the patient in supine position with the head backward (42.3%).

Regarding the attitude toward the first aid management of epistaxis. The majority (39.4%) of the respondents demonstrate the correct position which is holding the head forward rather than backward. 61.5% of respondents demonstrated the correct site for pinching the nose (soft part), While 38.5% of the respondents demonstrated the incorrect site (bony part). The main source of the respondents' knowledge regarding first aid management of epistaxis was Medical Curriculum (31.2%) followed by the Self-taught (23.2%).

Conclusion: The level of Knowledge and attitude regarding first aid management of epistaxis was good among most of the students. Health related specialties students have adequate knowledge on the standard first aid measures of epistaxis.

Keywords: Epistaxis, First Aid, Health Related Specialties Students.

*Correspondence to:

Dr. Muhannad A. Asiri,
MBBS, Intern, College of Medicine,
King Khalid University, Abha, Saudi Arabia.

Article History:

Received: 02-10-2017, **Revised:** 29-10-2017, **Accepted:** 18-11-2017

Access this article online	
Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2017.3.6.046	

INTRODUCTION

Epistaxis is the most common ear, nose and throat complaint presenting to an Emergency department. Epistaxis is a bleeding from inside the nose or nasal cavity. The vast majority of nose bleeds (90%) arise from Little's area on the anterior part of the nasal septum (Rains & Ritchie, 1984).¹ It is estimated from a Scandinavian survey in 1974 of 410 people that up to 60% of the population will experience one episode of epistaxis in their lifetime and 6% will seek medical attention.^{2,3} Other Reports have

mentioned an incidence ranging between 10%-60% of populations who have suffered at least one significant episode in their life time.⁴ It is clearly evident that the problem of epistaxis constitutes a significant amount to the workload in accident and emergency as well as otolaryngology departments.

Whilst some epistaxis episodes may do require an active intervention and even hospital admission, A vast majority of these patients settle with simple standard first aid Measures.⁵ The basic

first aid management of epistaxis is clearly mentioned in the guidelines. However, many surveys have suggested that these principles are not understood by patients and are not being well conveyed to patients by their doctors.⁶

To certain degree, health related specialties students are taught how to handle emergencies in a hospital emergency setting where drugs and other necessities are available. However, the adequate knowledge required for handling an emergency without hospital setting at the site of the accident or emergency may not be sufficient.^{7,8}

Although many studies have been written on the treatment of epistaxis, the knowledge and attitude of health related specialties students on this subject have not been documented. The purpose of this study was thus to assess and to promote the adequate knowledge and attitude regarding first aid management of epistaxis among health related specialties students at king Khalid university, Abha city, Saudi Arabia.

MATERIALS AND METHODS

A cross-sectional study was carried out amongst students of medical students at King Khalid University KKU (Dentistry College, medicine colleges, Pharmacy College, applied medical sciences college). Ethics approval, Anonymity of respondents, consent form, and other ethical issues were considered. The objectives of this study were to assess knowledge and attitude regarding first aid management of epistaxis among medical specialties students at King Khalid University, Abha, Saudi Arabia. Students start from level three enrolled (second academic year) were eligible for the study.

The purpose of the study was explained to the participants. The data was collected by using an electronic, semi-structured questionnaire. Which included questions on personal data (age, gender, GPA, specialty, clinical year and residency). Also consistent of 2 section to assess the knowledge and attitude of the students regarding the epistaxis and its first aid management. Data were tabulated by using Microsoft office Excel sheet, entered and analyzed by using SPSS, version 20.0. Ethical Committee approval was obtained before starting the study. The Chi-square test was used to find out the statistical significance of the differences in the proportions.

RESULTS

Out of 314 students participated in this study, 205 (65.3%) were male and 109 (34.7%) were female. medical students were the most respondents (52.2%) while (47.8%) were other medical specialties. Majority of the respondents were fourth year students. (Table 1) Out of 314 students; 164 (52.2%) were from medicine college, 82 (26.1%) from dental college, 37 (11.7%) from pharmacy college and 31 (9.8%) were from applied medical college. The mean age was 21.72±1.56 years (range, 18–26 years). 52 (16.4%) were in second year (level 3 and 4), 62 (19.7%) were in third year (level 5 and 6), 102 (32.4%) in fourth year (level 7 and 8), 49 (15.5%) in fifth year (level 9 and 10) and 49 (15.5%) were in sixth year (level 11 and 12). Their mean GPA was 3.9 with standard deviation 0.63. And 131 (41.7%) of students were received a first aid provider certificate, 183 (58.2%) had not received a first aid course. (Table 1)

Regarding the Knowledge of epistaxis, (239; 76.1%) of the respondents think that epistaxis is an emergent case. Regarding

the etiology, (98; 31.2%) of the participants respond that Bleeding Disorder is the commonest cause. Next common cause was Hypertension (57; 18.1%), followed by Injury to nose, including finger nail trauma (42; 13.3%). (Table 2)

The results regarding the knowledge among participants for seeking medical care during the attack Shows that 266 (84.7%) respondents said that a nosebleed cannot be stopped after 10 to 20 minutes of direct pressure is one of the commonest cause to seek for emergency care followed by a nosebleeds become more severe or more frequent 210 (66.8%). (Table 3)

The commonest first aid measures reported to be known by respondents when the patients get shocked ,were Pinching the nose (86.3%), Nasal packing (78.9%), Putting the patient in supine position with the head lowered (59.8%), giving Anti shock treatment (56.6%) and Putting the patient in supine position with the head backward (42.3%). (Table 4)

Regarding the attitude toward the first aid management of epistaxis The majority 39.4% of the respondents demonstrate the correct position which is holding the head forward rather than backward and 26.7% overall gave the correct duration of pinching the nose. (Table 5)

61.5% of respondents demonstrated the correct site for pinching the nose picture (A), While 38.5% of the respondents demonstrated the incorrect site picture (B). (Figure1)

The main source of the respondents' knowledge regarding first aid management of epistaxis was Medical Curriculum (31.2%) followed by the Self-taught (23.2%). (Table 6)

Table 1: Sociodemographic data

	N	%
GENDER		
Male	205	65.3%
Female	109	34.7%
COLLEGE		
Medicine	164	52.2%
Dentistry	82	26.1%
Pharmacology	37	11.7%
Applied Medical Sciences	31	9.8%
LEVEL OF EDUCATION		
Level 3 (2st year)	31	9.8%
Level 4 (2nd year)	21	6.6%
Level 5 (3rd year)	29	9.2%
Level 6 (3th year)	33	10.5%
Level 7 (4th year)	46	14.6%
Level 8 (4th year)	56	17.8%
Level 9 (5th year)	25	7.9%
Level 10 (5th year)	24	7.6%
Level 11 (6th year)	31	9.8%
Level 12 (6th year)	18	5.7%
AGE IN YEARS (Mean ±SD)	21.7±1.56	
GPA SCORE (Mean ±SD)	3.9±0.63	
RECEIVE A FIRST AID PROVIDER CERTIFICATE		
Yes	131	41.7%
No	183	58.2%

Table 2: Causes of epistaxis

	N	%
Epistaxis is considered as one of the emergency situations:		
Yes	239	76.1%
No	75	23.9%
Most common cause of Epistaxis :		
Bleeding disorders	98	31.2%
Hypertension	57	18.1%
Irritation due to allergies, colds, sneezing or sinus problems	26	8.2%
Nasal fracture	23	7.3%
Injury to nose, including finger nail trauma	42	13.3%
Deviated septum	18	5.7%
Overuse of decongestant nasal sprays	26	8.2%
Medications including analgesics and anticoagulant	24	7.6%

Table 3: When to seek for emergency care

When to seek for emergency care	N	%
A nosebleed cannot be stopped after 10 to 20 minutes of direct pressure.	266	84.7%
Nosebleeds recur 4 or more times in 1 week after you have tried Prevention measures.	210	66.8%
Nosebleeds become more severe or more frequent.	241	76.7%
After a head trauma	172	54.7%

Table 4: What should you do if the patient gets shocked

What should you do if the patient get shocked	N	%
Pinching the nose	271	86.3%
Nasal packing	248	78.9 %
Put the patient in supine position with the head lowered.	188	59.8%
Put the patient in supine position with the head backward.	133	42.3%
Examine the nose	46	14.6%

Table 5: Most proper position that patient with epistaxis

Most proper position that patient with epistaxis	N	%
Sitting with head tilted forward	124	39.4%
Sitting with head tilted backward	84	26.7%
Lying down and elevate the legs	31	9.8%
Lying down with ice pack over the nasal bridge	63	20%
Don't know	12	3.8%

Table 6: Sources of the respondents' knowledge

Sources of the respondents' knowledge	N	%
Medical curriculum	98	31.2
Self-taught	73	23.2
First aid course	61	19.4
Media	48	15.3
Guessing	34	10.8



Figure 1: The site of pinching the nose.

DISCUSSION

Epistaxis is the most common nasal emergency which defined as acute hemorrhage from the nostril or nasal cavity. It is a frequent emergency condition presented to emergency department. However, the vast majority of patients who present to the ED with epistaxis may treated successfully by an emergency physician by performing first aid management.⁹

First aid provider should be able to assess and provide appropriate medical care.

So our present study aimed to assess and to promote the adequate knowledge and attitude regarding first aid management of epistaxis among health related specialties students in king Khalid university, Abha, Saudi Arabia 2017 .

The respondents in this study were students from different medical specialties including Medicine, dentistry, pharmacology and Applied Medical Sciences.

It was expected that their levels of education will positively influence the knowledge and the attitude on the first aid management of epistaxis .

Regarding the Knowledge of epistaxis, 75.7% of the respondents think that epistaxis is an emergent case and 31.2% of them respond that Bleeding Disorder is the commonest cause of epistaxis.

The commonest first aid measures reported to be known by respondents when the patients get shocked ,were Pinching the nose (86.3%), Nasal packing (78.9%), Putting the patient in supine position with the head lowered (59.8%), giving Anti shock treatment (56.6%) and Putting the patient in supine position with the head backward (42.3%).

These results were not far from the result of study conducted by P. Mugwe in 2014¹⁰, in which the first aid measure known by most of the respondents was pinching the nose (94.0%) and nasal packing (80.6%). Unlike Several studies conducted by Adhikari in 2006¹¹, Ho EC 2008¹², and Klossek 2006¹³, found nasal packing to be the most common first line measure used by emergency clinical staff.

This may be attributed to lack of adequate knowledge on the first aid measures and lack of training in first aid in previously mentioned studies.

In general, the Knowledge and attitude towards first aid in epistaxis was good, The results of our study showed that The majority (39.4%) of the respondents demonstrate the correct position which is holding the head forward rather than backward and (26.7%) overall gave the correct duration of pinching the nose The Results Similar to the results of P. Mugwe in 2014¹⁰ in which Sixty percent (60%) of respondents described the correct position which a patient with nose bleeding should be placed.

In contrast to study done by Strachan¹⁹ only 36% gave a correct position. Regarding the correct site for pinching the nose. Only 44.3 % of respondents demonstrated the correct site, While the majority 55.7% of the respondents demonstrated the incorrect site. In accordance to P. Mugwe in 2014¹⁰ only 38.1% correctly demonstrated pinching the nose at the alae nasi. In general, the attitude of the medical specialties students towards first aid in epistaxis was good.

CONCLUSION

The level of Knowledge and attitude regarding first aid management of epistaxis was good among most of the students, and this study shows health related specialties students have adequate knowledge on the standard first aid measures of epistaxis and good attitude to provided first aid to patients presenting with epistaxis.

REFERENCES

1. Murray JAM: Epistaxis. In: Maran AGD, ed. Logan Tunner's Diseases of the Ear, Throat and Nose. 10th ed., John Wright, Brussels. 1988; 30: 234-238.
2. Santos PM, Lepore ML. Epistaxis. In: Bailey BJ, Calhoun KH (Eds). Head and Neck Surgery - otolaryngology. Lippincott-Raven, Philadelphia.1998; 513-529
3. Varshney S, Saxena RK. Epistaxis: a retrospective clinical study.Indian Journal of Otolaryngology, Head Neck Surgery. 2005; 57:125-129
4. Petruson B, Rudin R. The frequency of epistaxis in a male population sample. Rhinology1975;13(3):129-33.
5. Mcgarry G. W., Moulton C. The first aid management of epistaxis by Accident and Emergency department staff. Archives of Emergency Medicine. 1993; 10: 298-300.
6. Tanner R, HarneyMS.The initial management of epistaxis. Ir Med J. 2015 Apr;108(4):123-4 .
7. Khan A, Shaikh S, Shuaib F, Sattar A, Samani SA, Shabbir Q, et al. Knowledge attitude and practices of undergraduate students regarding first aid measures. J Pak Med Assoc 2010;60:68-72 .
8. Tekian A. Have newly graduated physicians mastered essential clinical skills? Med Educ 2002;36:406-7 .
9. Van Wyk FC, Massey S, Worley G, Brady S. Do all epistaxis patients with a nasal pack need admission? A retrospective study of 116 patients managed in accident and emergency according to a peer reviewed protocol. J Laryngol Otol. 2007 Mar. 121(3):222-7
10. P. Mugwe¹, K. J. Kamau²and O.K. Nyambaka³, Knowledge, Attitude and Practice in First Aid Management of Epistaxis by Accident and Emergency Clinical Staff at Kenyatta National Hospital, COSECSA/ASEA Publication -East and Central African Journal of Surgery. March/April 2014 Volume 19 (1)
11. Adhikari P, NM Thapa, BK Sinha. Aetiology and management of epistaxis at TU TeachingHospital. Journal of Institute of Medicine. 2006; 28(2): 23-44.
12. Ho EC, Chan JY. Front-line epistaxis management: let's not forget the basics. J Laryngol Otol. 2008; 122(7):696-9.
13. Klossek JM et al. Epistaxis and its management: an observational pilot study carried out in 23 hospital centres in France. Rhinology. 2006; 44(2):151-5.

Source of Support: Nil. **Conflict of Interest:** None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Muhannad A. Asiri, Sarah M. Mujarri, Shahad S. Alkhashrami, Amal A. Alghamdi, Khalid A. Almalwi, Saeed M. Asiri. Knowledge and Attitude Regarding First Aid Management of Epistaxis Among Students of Medical Colleges at King Khalid University, Abha City, Saudi Arabia. Int J Med Res Prof. 2017 Nov; 3(6):237-40. DOI:10.21276/ijmrp.2017.3.6.046