

Usefulness of Ultrasonography with Fine Needle Aspiration Cytology in Early Detection of Breast Cancer: A Hospital Based Study

Dinesh Kumar¹, Poonam Burdak^{2*}, Jitendra Acharya³

¹Assistant Professor, Department of Radiodiagnosis, S.P. Medical College, Bikaner Rajasthan, India.

²Assistant Professor, Department of Pathology, DDU Medical College, Churu, Rajasthan, India.

³Senior Demonstrator, Department of Dentistry, S.P. Medical College, Bikaner Rajasthan, India.

ABSTRACT

Introduction: Breast malignancy is the mainly extensively recognized tumor in females around the world. It is likewise the guideline reason for death from malignancy among ladies internationally. The (USG-FNAC) ultrasonography-fine needle yearning cytology procedures have adequate symptomatic legitimacy files and can be utilized for early arranging of bosom disease patients. The present examination was directed to evaluate breast malignancy utilizing FNAC and USG among consider population.

Materials and Methods: This study was conducted in the department of Radiodiagnosis in year 2017 at P. B. M. Hospital & S. P. Medical College. Bikaner. It included 80 women with breast lump. All were subjected to ultrasound, FNAC and incisional biopsy. The FNAC was performed using a 23G needle and 10 ml syringe with an average of four to six passes with constant suction.

Results: Out of 80 patients, 32 had education upto high school and 48 were illiterate. 74% had lump within 2- 6 months and 26% had more than 6 months. 54% had fever, 22% had fever, 24% had discharge from nipple, 5% had redness over breast and 3% had pain in opposite breast. Benign growth was seen in 50 (62%) patients and malignant growth was seen in 30 (38%) patients. Among benign, 90% was localized and 10% was ulcerated. In malignant, 55% was localized, 30% was ulcerated and 20% was infected fungating growth. USG had Sensitivity of 91%, specificity of 92% and PPV of 74.2%

whereas FNAC had Sensitivity of 95%, specificity of 98% and PPV of 94%. Various histopathological lesions were fibroadenoma (19), fibrocystic disease (17), DCIS (6), fat necrosis (3), tuberculous mastitis (2), adenosis (1), tubular adenoma (2) and breast cancer (30).

Conclusion: Early detection of breast lesions using fine needle aspiration cytology and ultrasonography is useful measure. Breast lump more than 1 month should be considered for detection and delay should be avoided as the chances of breast cancer cannot be over ruled.

Keywords: Fine Needle Aspiration Cytology, Fibroadenoma, Ultrasonography.

*Correspondence to:

Dr. Poonam Burdak,
Assistant Professor,
Department of Pathology,
DDU Medical College, Churu, Rajasthan, India.

Article History:

Received: 27-07-2018, Revised: 22-08-2018, Accepted: 29-09-2018

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2018.4.5.067	

INTRODUCTION

Breast cancer is the most well-known growth in females around the world. It is additionally the guideline reason for death from malignancy among ladies all around. The expression "bosom disease" alludes to a harmful tumor that has grown for the most part from the ductal epithelial cells in the breast.¹

As of now, it is the most driving reason for malignancy passing with 198,000 passings for every annum which speaks to 15.4% of all passings in created districts after that of the lung disease. In creating nations, it is the main driving reason for death among ladies with 324,000 passings which spoke to 14.3% of all passings. Moreover, this rate varied from 6 to 20 per 100,000 in East Asia and West Africa as a Whole.² The most common risk factors of breast cancer are obesity, lack of exercise, alcohol

consumption, smoking and use of oral contraceptive pills. Whereas genetic mutations and family history of breast cancer also plays important role. In general, women from higher socio-economical groups, regardless of their race/ethnicity, have higher rates of occurrence of breast cancer. Most of the breast cancers are asymptomatic.³

At present, screening for breast tumor in females is viewed as powerful approach to decrease rate of bosom malignancy. So a quick and precise conclusion device is vital and essential in the execution of bosom malignancy screening. The USG-FNAC ie ultrasonography-fine needle yearning cytology methods have satisfactory analytic legitimacy lists and can be utilized for early organizing of bosom malignancy patients.⁴

MATERIALS AND METHODS

This study was conducted in the department of radiodiagnosis in year 2017 at P.B.M. Hospital Bikaner Rajasthan. It included 80 women with breast lump. All were informed regarding the study and written consent was obtained. Patient information such as name, age, duration of lump etc. was recorded. A thorough history and physical examination was done in all patients.

All were subjected to ultrasound, FNAC and incisional biopsy. The FNAC was performed using a 23G needle and 10 ml syringe with an average of four to six passes with constant suction. We received tissue of breast lumps excised in Surgery OT under local/general anesthesia. Where the lump was very big, incision biopsy was done at proper site and with proper orientation of the incision. The histopathology reports gotten at follow-up visits were utilized to look at the consequence of USG and FNAC. High

recurrence ultra sound picture was utilized for analysis. Strong sores with sporadic shape, unpredictable edge, non-uniform appropriation of inside echotexture, shadowing and calcifications in hypo-echoic injuries are suggestive of growth by USG.

Table 1: Demographic parameters in patients

Total patients = 80	
Literacy	
Up to high school	32
Illiterate	48
Time lapse since lump first noticed	
2-6 months	74%
>6 months	26%

Table 2: Ultrasonography and FNAC in breast cancer

Test	Biopsy+/ breast cancer	Biopsy-/ no of breast cancer	Total	Sensitivity, specificity and PPV
USG				
Benign	2	45	47	Sensitivity 91%, specificity 92% and PPV 74.2%
Malignant	26	7	33	
FNAC				
Benign	1	48	49	Sensitivity 95%, specificity 98% and PPV 94%
Malignant	25	6	31	

RESULTS

Table 1 shows that out of 80 patients, 32 had education upto high school and 48 were illiterate. 74% had lump within 2- 6 months and 26% had more than 6 months. Figure 1 shows that 54% had fever, 22% had fever, 24% had discharge from nipple, 5% had redness over breast and 3% had pain in opposite breast.

Figure 2 shows that benign growth was seen in 50 (62%) patients and malignant growth was seen in 30 (38%) patients. Among benign, 90% was localized and 10% was ulcerated. In malignant, 55% was localized, 30% was ulcerated and 20% was infected fungating growth. Table 2 shows that USG had Sensitivity of 91%, specificity of 92% and PPV of 74.2% whereas FNAC had Sensitivity of 95%, specificity of 98% and PPV of 94%.

DISCUSSION

Breast cancer is recognised as a major public health problem in developing countries. Breast cancer is cancer that develops from breast tissue. Signs of breast cancer may include a lump in the breast, a change in breast shape, dimpling of the skin, fluid coming from the nipple, or a red scaly patch of skin. In those with distant spread of the disease, there may be bone pain, swollen lymph nodes, shortness of breath, or yellow skin.⁵

The two most commonly used screening methods, physical examination of the breasts by a healthcare provider and mammography, can offer an approximate likelihood that a lump is cancer, and may also detect some other lesions, such as a simple cyst. When these examinations are inconclusive, fine needle aspiration, or fine needle aspiration and cytology— FNAC) procedure is helpful in establishing the diagnosis. The needle

aspiration may be performed in a healthcare provider's office or clinic using local anaesthetic if required. A finding of clear fluid makes the lump highly unlikely to be cancerous, but bloody fluid may be sent off for inspection under a microscope for cancerous cells. Together, physical examination of the breasts, mammography, and FNAC can be used to diagnose breast cancer with a good degree of accuracy. USG is useful in detection of breast masses.⁶

In present study we analyzed patients with breast masses using FNAC and USG. Out of 80 patients, 32 had education upto high school and 48 were illiterate. 74% had lump within 2- 6 months and 26% had more than 6 months. We found that common symptoms in patients were fever, discharge from nipple, redness, over breast and pain in opposite breast. This is in accordance to Vissa et al.⁷

We found that among study population, benign growth was seen in 62% patients and malignant growth was seen in 38% patients. Among benign, 90% was localized and 10% was ulcerated. In malignant, 55% was localized, 30% was ulcerated and 20% was infected fungating growth. A study conducted by Nair MK⁸ found 71.2% of benign lesions.

A study conducted by S Usmani⁹ reported high specificity and sensitivity of FNAC as compared to USG. In our study, USG had Sensitivity of 91%, specificity of 92% and PPV of 74.2% whereas FNAC had Sensitivity of 95%, specificity of 98% and PPV of 94%.

In our study, histopathological lesions were fibroadenoma, fibrocystic disease, DCIS, fat necrosis, tuberculous mastitis, adenosis, tubular adenoma and breast cancer. This is in accordance to Y singh et al.¹⁰

CONCLUSION

Early detection of breast lesions using fine needle aspiration cytology and ultrasonography is useful measure. Breast lump more than 1 month should be considered for detection and delay should be avoided as the chances of breast cancer cannot be over ruled.

REFERENCES

1. F. Brunicaardi, Dana Andersen, Timothy Billiar, David Dunn, John Hunter, Jeffrey Matthews, Raphael E. Pollock. Schwartz's Principles of Surgery. 9th Edition. McGrawHill; 2010.
2. Xi-Wen Wang and, Yun-Hui Xiong and, Xiao-Qing Zen, Hai-Bo Lin, Qing-Yi Liu. Diagnostic Accuracy of Ultrasonograph Guided Fine-needle Aspiration Cytologic in Staging of Axillary Lymph Node Metastasis in Breast Cancer Patients: a Meta-analysis. *Asian Pacific J Cancer Prev.* 2001;13:5517-5523.
3. Gordon NH, Crowe JP, Brumberg DJ, Berger NA. Socioeconomic factors and race in breast cancer recurrence and survival. *Am J Epidemiol.* 1992;135:609-618.
4. Berg JW, Ross R, Latourette HB. Economic status and survival of cancer patients. *Cancer.* 1977;39:467-477.
5. Dayal HH, Power RN, Chiu C. Race and socio-economic status in survival from breast cancer. *J Chronic Dis.* 1982; 35:675-683.
6. Freeman HP, Wasfie TJ. Cancer of the breast in poor black women. *Cancer.* 1989;63:2562-2569.
7. Vissa Shanthi, Kashmir Ali, Nandam Mohan Rao, Baddukonda Appala, Rama Krishna, Kuppali Venkata Murali mohan. Clinicopathological study of breast lesions in females with assessment of correlation between tumor grade and prognostic factors. *J Biosci Tech.* 2011;367-378.
8. Nair MK, Sankaranarayanan R, Nair KS, Amma NS, V arghese C, Padmakumari G, et al. Overall survival from breast cancer in Kerala, India, in relation to menstrual, reproductive and clinical factors. *Cancer.* 2006;71:1791-6.
9. S Usmani, H Khan, N Ahmed, F Marafi, and N Garvie. Scintimammography in conjunction with ultrasonography for local breast cancer recurrence in post-mastectomy breast. *The British Journal of Radiology.* 2010;934-939.
10. Yumjaobabu Singh Takhellambam, Sunil Singh lourembam, opendro Singh Sapam, raju Singh KSheTrimaYum, bhupon Singh ningThoujam, TouSif Khan. Comparison of Ultrasonography and Fine Needle Aspiration Cytology in the Diagnosis of Malignant Breast Lesions. *JCDR.* 2013; 15-19.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Dinesh Kumar, Poonam Burdak, Jitendra Acharya. Usefulness of Ultrasonography with Fine Needle Aspiration Cytology in Early Detection of Breast Cancer: A Hospital Based Study. *Int J Med Res Prof.* 2018 Sept; 4(5):299-301. DOI:10.21276/ijmrp.2018.4.5.067