

A Relative Study on Laryngeal Mask Airway Lubrication with 0.005% Beclomethasone Cream v/a 2% Lidocaine

Bhaidas Onkar Patil¹, Suhasini Ravindra Sonavdekar^{2*}, Ranjan Mathur³

¹Associate Professor, Department of Anaesthesia,

SMBT Institute of Medical Sciences & Research Center, Dhamangaon Tal. Igatpuri, Nashik, Maharashtra, India.

^{2*}Associate Professor, Department of Anaesthesia,

MGM Medical College, Kamothe, Navi Mumbai, Maharashtra, India.

³Senior Professor and Head, Department of Dentistry, S. P. Medical College, Bikaner, Rajasthan, India.

ABSTRACT

Background: Postoperative sore throat (POST) is a slight complication after general anaesthesia. Numerous agents have been used as lubricant to lessen the incidence of POST with variable efficacy.

Methods: We compare the incidence of POST with 0.05% betamethasone gel and 2% lignocaine jelly as a lubricant for LMA insertion in patients undergoing general anaesthesia. Sixty subjects were separated randomly into two groups. Patients in Group B (n = 30) had 2.5 ml of 0.05% of betamethasone gel while the Group L had 2.5 ml of 2% lignocaine jelly applied on the cuff of LMA. After criterion induction and insertion of LMA cuff inflated to 60 cm of H₂O and was maintained at the same throughout the surgery.

Results: In PACU, patients evaluated about sore throat at immediate and 24 h post-operative period. POST was not seen in any of the patients of Group B. In group L 13% of the patients had 1st degree and 3% had 2nd degree of sore-throat in immediate post-operative period. After 24 h 3% patients had 1st degree sore-throat and 1% patients had 2nd degree of sore-throat in Group L patients.

Conclusion: We conclude that lubricating cuff of LMA with 0.05% of betamethasone gel is efficient in reducing the incidence of POST.

Keywords: Sore Throat; Betamethasone Gel; Lignocaine Jelly.

*Correspondence to:

Dr. Suhasini Ravindra Sonavdekar,

Associate Professor,

Department of Anaesthesia,

MGM Medical College,

Kamothe, Navi Mumbai, Maharashtra, India.

Article History:

Received: 05-04-2019, Revised: 01-05-2019, Accepted: 27-05-2019

Access this article online

Website: www.ijmrp.com	Quick Response code 
DOI: 10.21276/ijmrp.2019.5.3.045	

INTRODUCTION

Postoperative sore throat (POST) though a minor complication after general anaesthesia can be distressing to the patients. This is because of lack of airway humidity, trauma during airway insertion and suctioning, high anaesthetic air flow rates and surgical manipulation of airway and adjacent tissue. Many agents have been used as lubricant to reduce the incidence of postoperative sore throat with variable efficacy.^{1,2} We conducted a study to compare the incidence of postoperative sore throat with 0.05% betamethasone gel and 2% lignocaine jelly as a lubricant for laryngeal mask airway (LMA) insertion in patients undergoing general anaesthesia for elective surgery.

MATERIALS AND METHODS

It was a comparative study done on 60 patients of either sex, aged between 18–60 years (ASA physical status I or II) undergoing general anaesthesia. Patients requiring surgeries of expected

duration of more than 30 min and less than 2 h were included. Subjects were divided randomly in two groups by using sealed opaque envelopes, which were opened by the investigator just before the induction of anaesthesia. Assuming the proportion of sore throat with lignocaine as 0.5 and to test the specified difference in proportion of 0.4 with betamethasone group at 90% power the sample size comes out to be 30 patients in each group. It was a single blinded study. Only the subjects were blinded. After institutional ethical committee approval, written informed consent was obtained from all the patients. Patients having cervical spine disease, known airway obstruction, mouth opening less than 2.5 cm, any history of sensitivity to drugs i.e. lignocaine and betamethasone, with history of upper respiratory tract infections in the last 10 days, scheduled for oral, head and neck surgery, positioned in lateral or prone for surgery and patients who were at risk of aspiration were excluded from the study. Patients in group

B (n = 30) had 2.5 ml of 0.05% of betamethasone gel applied on the cuff of LMA. Patients in group L (n = 30) had 2.5 ml of 2% lignocaine jelly applied on the cuff of LMA.

In operating room, after the establishment of intravenous line and attachment of standard monitors, a standard intravenous anaesthetic induction technique involving fentanyl, thiopentone sodium and vecuronium bromide was used in all the patients. Maintenance of anaesthesia was carried out using 40% oxygen in 60 %medical air and isoflurane. LMA of appropriate size was introduced after lubrication of the surface with 2.5ml 0.05% betamethasone gel (Betagel, Microlabs Ltd.) or 2.5 ml 2% lignocaine jelly. LMA was introduced in deflated state with

introducer technique. Cuff was then inflated with air to a pressure equivalent to 60 cm of H₂O, using manometer. Proper placement of LMA was confirmed by capnograph. Intra-cuff pressure was monitored at 15, 30, 45, 60 and 90 min and just before extubation by using manometer. If cuff pressure was found to be increased, it was resettled to 60 cm of H₂O by deflating the cuff. At the end of surgery, patients were reversed with glycopyrrolate and neostigmine. In PACU patients were inquired about sore-throat at immediate and 24 h post-operatively and categorized as per the results were analysed using percentages, proportions and chi-square test. The statistical analysis was carried out using SPSS version 10.0.

Table 1: Degree of sore throat

0	No sore throat at any time since operation
1	Minimal sore throat (disappearing within 6 h)
2	Moderate sore throat (disappearing within 12 h)
3	Severe sore throat (lasting for 12 h or more)

Table 2: Demographic data and duration of surgery

	Group B (n=30)	Group L (n=30)
Age (years) (mean ± S.D)	32.12 ± 15	30.84 ± 8.87
Sex male/ female no. of patients	17/13	14 / 16
Weight (kg) (mean ± S.D)	52.20 ± 2.47	50.83 ± 3.31
Duration of surgery (min) (mean ± SD)	45.00 ± 6.95	46.00 ± 5.25

Table 3: Postoperative sore throat with 0.05% betamethasone gel and 2% lignocaine jelly

	Mild sore throat	Moderate sore throat
Immediate postop		
Group B (n = 30) 0.05% betamethasone gel	0	0
Group L (n = 30) 2% lignocaine jelly	13 (43.4)	5 (15.7)
p value	0.000	0.036
After 24 hrs		
Group B(n = 30) 0.05% betamethasone gel	0	0
Group L (n = 30) 2% lignocaine jelly	3 (10.0)	1 (3.6)
p value	p > 0.005	p > 0.005

RESULTS

The demographic data and length of surgery were analogous in both the groups [Table 2]. Postoperative sore-throat was not observed in any of the patients of group B. In group L, 33% of the patients had 1st degree and 10% had 2nd degree of sore throat in immediate post-operative period. After 24 h 16% patients had 1st degree sore-throat and 3% patients had 2nd degree of sore-throat in group L patients [Table 3].

DISCUSSION

In this study it was observed that the frequency of postoperative sore throat was significantly less when cuff of LMA was lubricated with betamethasone gel as compared with lignocaine jelly. The incidence of POST following tracheal intubation and classic LMA

has been observed to be 12– 60% and 7–34% respectively.⁵⁻⁷ LMA has gained popularity as a tool for airway management both in elective and emergency procedures for difficult airway.⁸⁻¹⁰ Various studies have described the incidence of sore throat up to 23% after use of LMA.¹¹Therefore there is a need to reduce POST following insertion of LMA. The issues related to POST have also been addressed in a recent editorial by Scuderi. He commented that sore throat is a lay description of pharyngitis which itself can have a variety of causes and may also include a variety of symptoms including laryngitis, hoarseness, cough or dysphagia.¹² The incidence of sore throat varies with use of different lubricants, degree of intra cuff pressures and number of attempts of airway device insertion.¹³Various lubricants have been used to reduce incidence of POST following LMA insertion with variable efficacy.

De Villager et al compared xylocaine 10% spray, silko spray, endos gel and water as lubricants for classic LMA and found that no single lubricant was associated with a statistically significant lower incidence of post-operative sore throat.¹⁴ McHardy and Chung in a review about POST commented that lubricants containing local anaesthetics do not appear to be beneficial but may actually be harmful. 2% lignocaine jelly has also been implicated as a cause of transient bilateral recurrent laryngeal nerve palsy.¹⁵ In a study comparing saline and lignocaine gel for occurrence of POST with classic LMA, the incidence of sore throat was found to be similar with both and author suggested that there is no role of lignocaine gel in prevention of post-operative sore throat.¹⁶ However Tanaka et al in a data based systematic review concluded that topical and systemic lidocaine is useful for prevention of post-operative sore throat.¹⁷ The efficacy of strepsil tablets, inhaled fluticasone propionate and benzydamine hydrochloride spraying on endotracheal tube cuff in reducing POST has also been evaluated recently.¹⁸⁻²¹ As there is a potential role of inflammation in causation of POST, the role of topical steroids in reducing POST has been studied by various workers. Others compared betamethasone gel and lignocaine jelly for lubrication of endotracheal tube and found it to be effective in reducing post-operative sore throat.²²⁻²⁴ Keeping in mind these studies, we used betamethasone gel for lubrication of LMA. Dosage of betamethasone, we used i.e. 6.25 mg (2.5 ml of 0.05%) was within the safe limits. Betamethasone gel has been used for lubrication of ETT without any adverse effects such as flaring up of local subtle infection. The role of intra cuff pressure in causing postoperative sore throat has been studied by some investigators, who found that a significant increase in cuff pressure is seen during first 60 min. There is no consensus whether intra cuff pressure of LMA contributes to POST. Burgard et al. evaluated the effect of laryngeal mask cuff pressure on incidence of POST and found that POST can be reduced when cuff pressure is continuously monitored and kept on low pressure values. Rieger et al. found that intra cuff pressure does not have any impact on POST following LMA insertion as high intra cuff pressure is unlikely to be transmitted directly to the mucosa; may be the elastic of the cuff stretches excessively and a proportion of the pressure measured is due to elastic recoil. However, we checked intra cuff pressure several times during surgery and kept it at 60 cm of H₂O. In conclusion, lubricating the cuff of LMA with betamethasone 0.05% gel may be effective in reducing the incidence of post-operative sore throat following elective surgeries. Since the sample size in our study was small, further studies involving a large number of patients are required to establish the role of betamethasone gel in preventing post-operative sore throat following LMA insertion.

CONCLUSION

We saw that lubricating cuff of LMA with 0.05% of betamethasone gel is efficient in reducing the incidence of POST.

REFERENCES

- Christensen AM, Willemoes-Larson H, Lundby L, Jakobsen KB. Postoperative throat complaints after tracheal intubation. *Brit J Anaesth* 1994;73:786-7.
- McHardy FE, Chung F. Postoperative sore throat: cause, prevention and treatment. *Anaesthesia* 1999;54:444-53.
- O'Connor Jr CJ, Borrromeo CJ, Stix MS. Assessing ProSeal laryngeal mask positioning: the suprasternal notch Test. *Anesth Analg* 2002;94(5):1374-5.
- Sumathi PA, Shenoy T, Ambareesha M, Krishna HM. Controlled comparison between betamethasone gel and lidocaine jelly applied over tracheal tube to reduce postoperative sore throat, cough and hoarseness of voice. *Brit J Anaesth* 2008;100:215-8.
- Mandoe H, Nikolajsen L, Lintrup U, Jepson D, Molgaard J. Sore throat after endotracheal intubation. *Anesth Analg* 1992;74:897-900.
- Joshi GP, Inagaki Y, White PF, Taylor-Kennedy L, Wat LI, Gevirtz C. Use of the laryngeal mask airway as an alternative to the tracheal tube during ambulatory anesthesia. *Anesth Analg* 1997;85:573-7.
- Dingle J, Whitehead MJ, Wareham K. A comparative study of the incidence of sore throat with the laryngeal mask airway. *Anaesthesia* 1994;49:251-4.
- Brimacombe J, Keller C, Fullekrug B, Agro F, Rosenbaltt W, Deirdorf SF, et al. A multicenter study comparing the ProSeal and classic laryngeal mask airway in anesthetized, nonparalyzed patients. *Anesthesiology* 2002;96(2):289-95.
- Natalini G, Lanza G, Rosano A, Dell Agnolo P, Bernardini A. Standard laryngeal mask airway and LMA-ProSeal during laparoscopic surgery. *J Clin Anaesth* 2003;15(6):428-32.
- Evans NR, Gardner SV, James MF, King JA, Roux P, Bennett P, et al. The ProSeal laryngeal mask: results of a descriptive trial with experience of 300 cases. *Brit J Anaesth* 2002;88(4):534-9.
- Woo YC, Cha SM, Kang H, Baek CW, Jung YH, Kim JY, et al. Less perilaryngeal gas leakage with SLIPA than PLMA in paralysed patients. *Can J Anaesth* 2011;58(1):48-54.
- Scuderi PE. Postoperative sore throat: more answers than questions. *Editorial Anesth Analg* 2010;111(4):831-2.
- Figueredo E, Martinez M, Pintanel T. A comparison of the ProSeal laryngeal mask and the laryngeal tube in spontaneously breathing anesthetized patients. *Anesth Analg* 2003;96:600-5.
- De Vlioger J, Soetens M, Van der Donck A. Sore throat with the laryngeal mask: does the lubricant matter? *Acta Anaesthesiol Belg* 1996;47(4):195-8. [15] Inomata S, Nishikawa T, Suga A, Yamashita S. Transient bilateral vocal cord paralysis after insertion of a laryngeal mask airway. *Anesthesiology* 1995;82:787-8.
- Keller C, Sparr HJ, Brimacombe JR. Laryngeal mask lubrication. A comparative study of saline versus 2% lignocaine gel with cuff pressure control. *Anaesthesia* 1997;52(6):592-7.
- Tanaka Y, Nakayama T, Nishimori M, Sato Y, Furuya H. Lidocaine for preventing postoperative sore throat. *Cochrane Database Syst Rev* 2009;8(3):CD004081.
- Ebneshaidi AE, Mohseni M. Strepsils tablets reduce sore throat and hoarseness after tracheal intubation. *Anesth Analg* 2010;111(4):892-4.
- Hung NK, Wu CT, Chan SM, Lu C, Huang YS, Yeh CC, et al. Effect on postoperative sore throat of spraying the endotracheal tube cuff with benzydamine hydrochloride, 10% lidocaine and 2% lidocaine. *Anesth Analg* 2010;111(4):882-6.
- Tazeh-kand NF, Eslami B, Mohammadian K. Inhaled flucatisone propionate reduces postoperative sore throat, cough and hoarseness. *Anesth Analg* 2010;111(4):895-8.

20. Huang YS, Hung NK, Lee MS, Kuo CP, Yu JC, Huang GS, et al. The effectiveness of benzydamine hydrochloride spraying on endotracheal tube cuff or oral mucosa for postoperative sore throat. *Anesth Analg* 2010;111(4):887–91.
21. Stride PC. Postoperative sore throat: topical hydrocortisone. *Anaesthesia* 1990;45:968–71.
22. Jeon YT, Na HS, Park SH, Oh AY, Park HP, Yun MJ, et al. Insertion of the ProSeal laryngeal mask airway is more successful with the 90 rotation technique. *Can J Anaesth* 2010;57(3):211–5.
23. Kazemi A, Amini A. The effect of betamethasone gel in reducing sore throat, cough and hoarseness after laryngotracheal intubation. *Middle East J Anaesthesiol* 2007;19(1):197–204.
24. Burgard G, Mollhoff T, Prien T. The effect of laryngeal mask cuff pressure on postoperative sore throat incidence. *J Clin Anesth* 1996;8:198–201. [26] Rieger A, Brunne B, Striebel HW. Intra cuff pressures do not predict laryngopharyngeal discomfort after use of the laryngeal mask airway. *Anesthesiology* 1997;87:63–7.

Source of Support: Nil.

Conflict of Interest: None Declared.

Copyright: © the author(s) and publisher. IJMRP is an official publication of Ibn Sina Academy of Medieval Medicine & Sciences, registered in 2001 under Indian Trusts Act, 1882.

This is an open access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Bhaidas Onkar Patil, Suhasini Ravindra Sonavdekar, Ranjan Mathur. A Relative Study on Laryngeal Mask Airway Lubrication with 0.005% Beclomethasone Cream v/a 2% Lidocaine. *Int J Med Res Prof.* 2019 May; 5(3):206-09. DOI:10.21276/ijmrp.2019.5.3.045